

ASTHMA HEALTH CARE PLAN

Child name:			Birth date:		
Community program name:					
Parent/guardian name:					
Home Ph#:	Cell #:		Work Ph#:		
Parent/guardian name:					
Home Ph#:	Cell #:		Work Ph#:		
Alternate emergency contact name:					
Home Ph#:	Cell #:		Work Ph#:		
Allergist:			Phone #:		
Pediatrician/Family doctor:			Phone #:		
Known allergies:					
Does child wear MedicAlert™ identification for asthma?					
TRIGGERS - List items that most commonly trigger your child's asthma.					
<u>RELIEVER MEDICATION</u> (or bronchodilators) provides fast temporary relief from asthma symptoms. It is recommended that Reliever medication is carried with the child so it is available if an asthma episode occurs.					
What Reliever medication has		☐ Salbutamol (e.g. Ventolin [®] , Airomir [®])			
prescribed for your child? (CH ONE)	ECK Symbico	☐ Symbicort [®] ☐ Oth			
How many puffs of Reliever	_ 1 puff _	1	or 2 puffs		
medication are prescribed for a asthma episode? (CHECK ONE	an 0	_	•		
astillia episode: (OIILON ONL		Ot	her		
Where does your child carry hi	s/her		herurse		
• `	s/her	ack p			
Where does your child carry hi	s/her fanny pa ONE) backpad	ack po	urse hertake their Relie		
Where does your child carry hi Reliever medication? (CHECK Does your child know when to	s/her	Can your child medication on	urse her take their Relie their own?	ever	
Where does your child carry his Reliever medication? (CHECK Does your child know when to their Reliever medication? CIRCLE the type of medication	s/her	Can your child medication on	urse her take their Relie their own?	ever	



ASTHMA HEALTH CARE PLAN

Name:	Birth date:		
IF YOU SEE THIS:	DO THIS:		
 Symptoms of asthma Coughing Wheezing Chest tightness Shortness of breath Increase in rate of breathing while at rest 	 Remove the child from triggers of asthma. Have the child sit down. Ensure the child takes Reliever medication (usually blue cap or bottom). Encourage slow deep breathing. Monitor the child for improvement of asthma symptoms. If Reliever medication has been given and asthma symptoms do not improve in 5-10 minutes, contact parent/guardian. Reliever medication can be repeated once at this time. If the child is not well enough to remain at the community program, the parent/guardian should come and pick them up. If any of the emergency situations occur (see list below), call 911/EMS. 		
 Emergency situations Skin pulling in under the ribs Skin being sucked in at the ribs or throat Greyish/bluish color in lips and nail beds Inability to speak in full sentences Shoulders held high, tight neck muscles Cannot stop coughing Difficulty walking 	 Activate 911/EMS. Delegate this task to another person. Do not leave the child alone. Continue to give Reliever medication as prescribed every five minutes. Notify the child's parent/guardian. Stay with the child until EMS personnel arrives. 		
Signs that asthma is not controlled	or wheezing.		
have reviewed this health care plan and provide conservations. Parent/guardian signature: have reviewed this health care plan to ensure it provide lurse signature:	Date:les the community program with required information.		
Pocumentation			

☐ Instruction sheet for medication device attached